FIG. 1

Constitutively Active Receptors

|               |                    | 1,  | /27                 |   |                                      |                                      |   |   | _ |
|---------------|--------------------|---|---------------------|---|--------------------------------------|--------------------------------------|---|---|---|
| Reference     |                    | (Robbins, Nadeau et al. 1993)                               |                     | (Pauwels, Gouble et al.<br>1999)                | (Egan, Herrick-Davis et<br>al. 1998) | (Herrick-Davis, Egan et<br>al. 1997) |   |   |   |
| Assay / Cells |                    | adenylyl cyclase activity/<br>HEK293, stably<br>transfected |                     | binding of [ <sup>15</sup> S]GTP[S] /<br>CHO-KI | IP production / COS-7                | PI hydrolysis / COS-7                |   |   |   |
| Sequence      |                    | 92<br>VSIVL <u>R</u> TTIIL<br>K                             |                     | 313<br>RERKA <u>T</u> KTLØI<br>K, R, Q          | 322<br>NBQKAÇKVLGI<br>K              | 312<br>Neddagkvlgi<br>L              | Ě |   |   |
| Mutation Site |                    | IMI   |                     | C-terminus of IC3                               | C-terminus of IC3                    | C terminus of IC3                    |   |   |   |
| Receptor      |                    | melanocyte-stimulating hormone MSH                          |                     | 5-hydroxytryptamine <sub>1B</sub>               | 5-hydroxytryptamine <sub>2A</sub>    | 5-hydroxytryptamine <sub>ac</sub>    |   | , |   |
| File Name     | CLASS A<br>GROUP I | MSHR_mouse  | CLASS A<br>GROUP II | 5H1B_human                                      | 5H2A_human                           | 2H2C_rat                             |   |   |   |

\*0.g

.

| ŦŢ.      |  |
|----------|--|
| O1       |  |
| O1       |  |
| Ō        |  |
| F. 1     |  |
| <u>F</u> |  |
| ₹ .      |  |
|          |  |
|          |  |
| į.       |  |
| n.       |  |
|          |  |
| N        |  |
|          |  |
| ÷        |  |

|                  |                     | (Scheer, Fanelli et al. 1997) |             |                                    | (Scheer, Costa et al. 2000)     |             | (Perez, Hwa et al. 1996)         |                    |                          | (Hwa, Gaivin et al. 1997) |                          | (Kjelsberg, Cotecchia et    | al. 1992)                                    | (Allen, Lefkowitz et al.<br>1991)   | (Ren, Kurose et al. 1993)     |                  | (Högger, Shockley et al. 1995) |                             | n (Liu, Blin et al. 1996)   |                |
|------------------|---------------------|-------------------------------|-------------|------------------------------------|---------------------------------|-------------|----------------------------------|--------------------|--------------------------|---------------------------|--------------------------|-----------------------------|--|-------------------------------------|-------------------------------|------------------|--------------------------------|-----------------------------|-----------------------------|----------------|
|                  |                     | IP / COS-7                    |             |                                    | IP / COS-7                      |             | IP / COS-1                       | £                  | arachidonic acid release | IP/COS-1                  |                          | PI/COS-7                    |  | PI hydrolysis /<br>rat fibroblast   | adenylyl cyclase              | mmondon / new 23 | PI / HEK(U293)                 |                             | IP production, inhibition   | COS-7          |
|                  |                     | 63<br>PAIVGNILVIL             | ١٨          | 142<br>CAISI <u>D</u> RYIGV<br>A   | 143<br>CAISIDRYIGV              | К           | 128<br>AVDVI <u>C</u> CTASI<br>P | • 6                | REKKAAKTLGI              | ą                         | 204<br>Brppyllpsslg<br>V | 293                         | SKEKKAAKT<br>X=19 different<br>substitutions | 288 293<br>KPSREKKAAKTLGI<br>K H I. | 1 111 2                       | X=F, A, C, E, K  | 360<br>Slvkekkaartls           | Ą                           | 390                         | 1-4 A inserted |
| :                |                     | TMDI                          |             | junction between TMDIII<br>and IC2 | junction between TMDIII and IC2 |             | TMIII                            | onshown and of 102 | cationayi cita oi too    | TMV                       |                          | C-terminal IC3              |  | C-terminus IC3                      | C-terminal IC3 loop           |                  | C-terminal IC3 loop junction   |                             | junction of IC3 and TMVI    |                |
| FIG. 1 (2 of 15) |                     | α <sub>18</sub> -adrenergic   | alpha 1B-AR |                                    | α <sub>18</sub> -adrenergic     | alpha 1B-AR | α <sub>is</sub> -adrenergic      |                    |                          |                           |                          | α <sub>18</sub> -adrenergic |  | α <sub>18</sub> -adrenergic         | α <sub>2</sub> C10-adrenergic | alpha-2AAR       | muscarinic Hm1                 | muscarinic acetylcholine M1 | muscarinic acetylcholine M2 |                |
| FIG. 1           | CLASS A<br>GROUP II | A1AB_human                    |             |                                    | A1AB_human                      |             | A1AB_human                       |                    |                          |                           |                          | A1AB_human                  |  | A1AB_human                          | A2AA_human                    |                  | ACM1_human                     |                             | ACM2-human                  |                |

FIG. 1 (3 of 15)

|                     | <del></del>                                     | T  | T  | 3/27   | ,  |   |   |                           |                                       |
|---------------------|---|--|--|--|--|---|---|---------------------------|---------------------------------------|
|                     | (Blüml, Mutschler et al. 1994)                  | (Burstein, Spalding et al. 1996)               | (Spalding, Burstein et al. 1998)             | (Spalding, Burstein et al. 1997)                   | (Mason, Moore et al.<br>1999)              | (Samama, Cotecchia et al. 1993);<br>(Lefkowitz, Cotecchia et al. 1993)        | (Charpentier, Jarvie et al.<br>1996)                          | (Cho, Taylor et al. 1996) | (Alewijnse, Timmerman<br>et al. 2000) |
|                     | IP/COS-7  | β-gal / NIH 3T3                                | β-gal; radioligand<br>binding / NIH-3T3      | β-gal; radioligand<br>binding / NIH-3T3            | adenylyl cyclase; agonist<br>binding / CHW | adenylyl cyclase<br>activation; agonist<br>binding affinity /<br>COS-7 or CHO | adenylyl cyclase; cAMP<br>accumulation / HEK293               | CAMP accumulation / COS-7 | cAMP producti n /<br>HEK-293          |
|                     | 507<br>TWLPYLWIT<br>S                           | chimera composed of m21-69 m577-445 m2 391-466 | ALLLA RIITW TPYNI MVLVST M L H C V S F       | 465<br>YNIMVLV <u>S</u> TFCDKCV<br>X=V,F,R,K,+more | 389<br>RKAFQGLLCCA<br>R                    | 266 272<br>FCLKEHKALKTLGI<br>SR K A   | 264 SFKMSEKRETKVLKT I K 288 from DIB receptor APDTSIKKETKVLKT | 286<br>FVCCWLPPFIL<br>A   | 115<br>FMISL <u>D</u> RYCAV<br>N,A    |
|                     | TMVI  | N-terminus to TMII TMVI                        | TMVI   | junction of TMVI and EC3                           |  | C-terminal IC3 loop   | carboxyl terminal IC3   | TMVI                      | IC                                    |
|                     | m3 muscarinic (rat) muscarinic acetylcholine M3 | m5 muscarinic<br>muscarinic acetylcholine M5   | m5 muscarinic<br>muscarinic acetylcholine M5 | m5 muscarinic<br>muscarinic acetylcholine M5       | β <sub>1</sub> -adrenergic                 | β <sub>2</sub> -adrenergic<br>beta-2AR  | dopamine<br>DIA   | dopamine<br>D1            | histamine $ m H_2$                    |
| CLASS A<br>GROUP II | ACM3_rat  | ACM5_human                                     | ACM5_human                                   | ACM5_human   | B1AR_human                                 | B2AR_human  | DADR_human  | DADR_human                | HH2R_rat                              |

۹.

FIG. 1 (4 of 15)

|               |                      | 4,  | /27                                      |   |   |                           |
|---------------|----------------------|---|--|---|---|---------------------------|
| Reference     |                      | (Rim and Oprian 1995)   | (Acharya and Karnik<br>1996)             | (Han, Smith et al. 1998)                | (Govardhan and Oprian<br>1994);<br>(Cohen, Yang et al. 1993)  | (Cohen, Yang et al. 1993) |
| Assay / Cells |                      | transducin;<br>phosphorylation by<br>rhodopsin kinase / COS   | transducin; radioligand<br>binding / COS | transducin, GTPyS<br>uptake / COS       | transducin; radioligand binding / COS   |                           |
| Sequence      |                      | 90 PMVLGGFTSTLY D 113 GCNLEGFFAT Q 292 296 MTIPAFFAKSAAIY E G, E, M 23Ala neutral a.a converted to carboxylate and competes with 113Glu for salt bridge with 236Lys | 134<br>VVLAIERYVVV<br>I,Q,S              | 257<br>RMVIIMVIAFL<br>Y,N<br>plus G113Q | 296 PAFFAKSAAIY G X=E,M natural mutants + 10 different a.a. substitutions disrupts critical salt bridge between 296Lys(TMVII) and 113Glu(TMIII) | 134<br>VVLAIERYVVV<br>Q   |
| Mutation Site |                      | TMII<br>TMIII   | TMIII                                    | TM6<br>plus TM3                         | TMVII   | 22                        |
| Receptor      | ÷.                   | opsin<br>rhodopsin  | opsin<br>rhodopsin                       | opsin<br>rhodopsin                      | opsin<br>rhodopsin  |                           |
| File Name     | CLASS A<br>GROUP III | OPSD_human  | OPSD_human                               | OPSD_human                              | OPSD_human  |                           |

FIG. 1 (5 of 15)

| (Matus-Leibovitch,<br>Nussenzveig et al. 1995)                                     |  |  |
|--|--|--|
| "Ca 2" efflux, [Ca 2"] / Xenopus oocytes; IP formation / AtT20, stably transfected |  |  |
| 335<br>Frkl <u>c</u> nckok<br>Stop   |  |  |
| carboxyl tail  |  |  |
| thyrotropin-releasing hormone carboxyl ta<br>TRH-R                                 |  |  |
| TRFR_mouse   |  |  |

\*\*\*

FIG. 1 (6 of 15)

| File Name           | Receptor                  | Mutation Site Sequence | Sequence           | Assay / Cells         | Reference                 |
|---------------------|---------------------------|------------------------|--------------------|-----------------------|---------------------------|
| CLASS A<br>GROUP IV | CLASS A<br>GROUP IV       |                        |                    |                       |                           |
| BRB2_human          | bradykinin B <sub>2</sub> | TMII                   | 113<br>AIISMNLY8SI | IP production / COS-7 | (Marie, Koch et al. 1999) |
|                     | B2 bradykinin<br>BK-2     | TMVI                   | 256                |                       |                           |
|                     |                           |                        | nut ii Çanes Çi    |                       |                           |
|                     |                           |                        |                    | ,                     |                           |
|                     |                           |                        |                    |                       |                           |

ë.

FIG. 1 (7 of 15)

| •                                 |
|-----------------------------------|
|                                   |
| TMIII                             |
|                                   |
|                                   |
| C-terminus of TM7                 |
| other multiple mutations          |
| formylmethionylleucylphenylal IC1 |
|                                   |
|                                   |
| ICZ                               |
|                                   |
|                                   |
| වු                                |
|                                   |
| TMVI                              |
|                                   |
| TIM6                              |
|                                   |
| TMVI                              |
|                                   |
| TM3                               |
|                                   |
| <u>5</u>                          |
| ···                               |

osser acatem

| FIG. 1 (8 of 15) | (8 of 15)                                      |                        |   |   |                                      |
|------------------|--|------------------------|---|---|--------------------------------------|
| PAFR_human       | platelet-activating factor (PAF)               | C-terminus of IC3      | 231<br>EVKRRALMNVCTVLAV<br>R                      | IP production / COS-7   | (Parent, Le Gouill et al.<br>1996)   |
| PAFR_human       | platelet-activating factor (PAF)               | TMIII                  | 100<br>CLFFINTYCSV<br>A                           | arachnidonate release, IP production, adenylyl cylease inhibition / CHO | (Ishii, Izumi et al. 1997)           |
| PE23_human       | prostaglandin E,,<br>EP3III<br>EP3IV           | C-terminal tail        | 360 FCQBEFWGN FCQMRKRRLREGEFWGN Tcruncated        | inhibition of adenylyl cyclase / CHO-K1                                 | (Jin, Mao et al. 1997)               |
| PE23_mouse       | prostaglandin E,<br>EP3                        | carboxyl-terminal tail | 336 KILLRKFCQIRDHI (3α) MANHL (3β) ↑truncated     | inhibition of adenylate<br>cyclase / CHO, stably<br>expressed           | (Hasegawa, Negishi et al.<br>1996)   |
| THRR_human       | thrombin                                       | EC2 loop               | 259 268<br>CHDVINETLLEGYYAYY<br>DLKD KOP I        | 43Ca 24 efflux, PI<br>hydrolysis, reporter gene<br>induction / COS-7    | (Nanevicz, Wang et al.<br>1996)      |
| TSHR_buman       | thyrotropin (TSHR) thyroid stimulating hormone | BC1                    | 486<br>YYNHALDWQTG<br>F,M<br>568<br>YAKUSICLPMD   | inositol phosphate<br>diacylglycerol cascade /<br>COS-7                 | (Parma, Van Sande et al.<br>1995)    |
| TSHR_human       | thyrotropin (TSHR) thyroid stimulating hormone | TMIII                  | ASBLSYYTLTV ASPLSYYTLTV A 672 YPLNSCANPFL         | adenylyl cyclase<br>activation / COS-7                                  | (Duprez, Parma et al.<br>1994)       |
| TSHR_human       | thyrotropin (TSHR)                             | TMV                    | Y<br>597<br>VAFVI <u>V</u> CCCHV<br>L             | cAMP formation /<br>COS-7 cells   | (Esapa, Duprez et al.<br>1999)       |
| TSHR_human       | thyrotropin (TSHR) thyroid stimulating hormone | TMVII                  | AIFT  | cAMP formation /<br>CHO cells   | (Russo, Wong et al. 1999)            |
| TSHR_human       | thyrotropin (TSHR) thyroid stimulating hormone | <u> </u>               | 613 621<br>VRNPQ <u>YNPGDKDTK</u> IAK<br>deletion | cAMP formation /<br>COS-7   | (Wonerow, Schoneberg et<br>al. 1998) |

FIG. 1 (9 of 15)

| TSHR_human | TSHR_human thyrotropin (TSHR) | IC3 / TMVI | 623 632<br>KDTKIAKRMAVLIFIDFICM | cAMP activation / | (Paschke, Tonacchera et al. 1994) |
|------------|-------------------------------|------------|---------------------------------|-------------------|-----------------------------------|
|            | thyroid stimulating hormone   |            | I V                             | -                 | •                                 |
| V2R human  | vasopressin V2                | IC2        | 136                             | cAMP formation /  | (Morin, Cotte et al. 1998)        |
| 1          |                               |            | LAMTLDRHRAI                     | COS-1             |                                   |
|            |                               |            | <b>4</b>                        |                   |                                   |

FIG. 1 (10 of 15)

|               |                    |  |                     |  | 1                                       | O/ | 27      | 7         |   |                                   |                                  |  |                                   |
|---------------|--------------------|--|---------------------|--|---|----|---------|-----------|---|-----------------------------------|----------------------------------|--|-----------------------------------|
| Reference     |                    | (Cohen, Thaw et al. 1997)                |                     | (Schipani, Jensen et al.<br>1997)                |   |    |         |           | (Tseng and Lin 1997)                                | (Hjorth, Orskov et al.<br>1998)   |                                  | (Gaudin, Maoret et al.<br>1998)<br>(Gaudin, Rouyer-Fessard<br>et al. 1998) |                                   |
| Assay / Cells |                    | adenylyl cyclase cAMP production / COS-1 |                     | cAMP accumulation /<br>COS-7                     |   |    |         |           | cAMP production / L293                              | cAMP accumulation /<br>COS-7      |                                  | cAMP production /<br>COS-7 or CHO  |                                   |
| Sequence      |                    |  |                     | 223<br>TrnyiHmhlfl<br>R, K                       | 410<br>KLLKS <u>T</u> LVLMP<br>C,others |    |         |           | 340<br>VPAPV <u>T</u> EBQAR<br>P                    | 178<br>Trny i <u>H</u> gnlfa<br>R | 352<br>RLARS <u>T</u> UTLIP<br>A | 178 RNYIHMHLFI R requires functional integrity of the N-terminal BC domain | 343<br>LARSILLIP<br>X= K, P       |
| Mutation Site |                    | wild type (native) protein               |                     | junction of IC1 and TMII                         | junction of IC3 and TMVI                |    |         |           | TMVI  | junction of IC loop1 and<br>TMII  | IC end of TMVI                   | junction of IC loop 1 and TMII   | junction of IC loop 3 and<br>TMVI |
| Receptor      |                    | human calcitonin hCTR-1<br>hCTR-2        |                     | parathyroid hormone<br>PTH / PTH-related peptide |   |    |         |           | glucose-dependent<br>insulinotropic peptide (GIP-R) | glucagon                          |                                  | vasoactive intestinal peptide 1<br>(VIP)                                   |                                   |
| File Name     | CLASS B<br>GROUP I | CALR_human                               | CLASS B<br>GROUP II | PTRR_human                                       |   |    | CLASS B | GROUP III | GIPR_human  | GLR_rat                           |                                  | VIPR_human   |                                   |

FIG. 1 (11 of 15)

| File Name  | Receptor                   | Mutation Site | Sequence  | Assay / Cells | Reference                      |
|------------|----------------------------|---------------|---|---------------|--------------------------------|
| CLASS C    |                            |               |   |               |                                |
| CASR_human | CASR_human calcium-sensing | N-terminal BC | TLSFVAONKIDSIANIDEFCNCSEHI IP/tsA various substitutions, in multiple combinations | IP/tsA        | (Jensen, Spalding et al. 2000) |
|            |                            |               |   |               |                                |
|            |                            |               |   |               |                                |
|            |                            |               |   |               |                                |

FIG. 1 (12 of 15)

|               |         |                                |             |                           | 1                          | 2                        | /2             | 27       | <b>.</b> |                          |   |                  |                         |                            |                  |                         |                  |  |  |
|---------------|---------|--------------------------------|-------------|---------------------------|----------------------------|--------------------------|----------------|----------|----------|--------------------------|---|------------------|-------------------------|----------------------------|------------------|-------------------------|------------------|--|--|
| Reference     |         | (Olesnicky, Brown et al. 1999) |             | (Konopka, Margarit et al. | 1996)                      | (Dube, DeCostanzo et al. | 2000)          | •        |          |                          |   |                  |                         | (Boone, Davis et al. 1993) |                  | (Sommers, Martin et al. | 2000)            |  |  |
| Assay / Cells |         | heterologous yeast assay       |             | lacZ reporter gene        |                            | lacZ reporter gene /     | yeast          |          |          |                          |   |                  |                         | β-galactosidase            |                  | β-galactosidase         |                  |  |  |
| Sequence      |         | 229<br>Plsay <u>o</u> iyigt    | Ωŧ          | 258                       | QSLLV <u>PS</u> IIFI<br>LL | 223                      | MSFVLYVKLILAIR | <b>ာ</b> | 247 251  | DSFHI <u>LLIMS</u> COSLL | ຍ | double mutations | shaded double mutations | 194                        | DVRDILHCTNS<br>O | 253 258                 | LIMSCOSLLVPSIIFI |  |  |
| Mutation Site |         | TM6                            |             | TM6                       |                            | double mutations TM5     |                | and      |          | TM6                      |   |                  |                         | IC3                        |                  | TM6                     |                  |  |  |
| Receptor      |         | pheromone                      |             | pheromone α-factor        |                            | pheromone α-factor       |                |          |          |                          |   |                  |                         | pheromone a-factor         |                  | pheromone α-factor      |                  |  |  |
| File Name     | CLASS D | O74283<br>RCB2                 | C. cinereus | STE2_yeast                |                            | STE2 yeast               | 1              |          |          |                          |   |                  |                         | STE3_yeast                 |                  | STE2_yeast              | •                |  |  |

## FIG. 1 (13 of 15) Bibliography

Alewijnse, A. E., H. Timmerman, et al. (2000). "The Effect of Mutations in the DRY Motif on the Constitutive Activity and Structural Instability of the Histamine H(2) Receptor." Acharya, S. and S. S. Karnik (1996). "Modulation of GDP release from transducin by the conserved Glu 134-Arg 135 sequence in rhodopsin." LBiol Chem 271(41): 25406-11. Mol Pharmacol 57(5): 890-898.

Allen, L. F., R. J. Lefkowitz, et al. (1991). "G-protein-coupled receptor genes as protooncogenes: constitutively activating mutation of the alpha 1B-adrenergic receptor enhances mitogenesis and tumorigenicity." Proc Natl Acad Sci U S A 88(24): 11354-8.

Amairuda, T. T., 3rd, S. Dragas-Graonic, et al. (1995). "Signal transduction by the formyl peptide receptor. Studies using chimeric receptors and site-directed mutagenesis define a novel domain for interaction with G-proteins." J. Biol Chem 270(47): 28010-3.

Blüml, K., E. Mutschler, et al. (1994). "Functional role in ligand binding and receptor activation of an asparagine residue present in the sixth transmembrane domain of all muscarinic acetylch line receptors." I Biol Chem 269(29): 18870-6.

Boone, C., N. G. Davis, et al. (1993). "Mutations that after the third cytoplasmic loop of the a-factor receptor lead to a constitutive and hypersensitive phenotype." Proc Natl Acad

Sci USA 90(21): 9921-5.
Bradbury, F. A., N. Kawate, et al. (1997). "Post-translational processing in the Golgi plays a critical role in the trafficking of the luteinizing hormone/human chorionic gonadotropin receptor to the cell surface." LBiol Chem 272(9): 5921-6.
Bradbury, F. A. and K. M. Menon (1999). "Evidence that constitutively active luteinizing hormone/human chorionic gonadotropin receptors are rapidly internalized."

13/27

Burger, M., J. A. Burger, et al. (1999). "Point mutation causing constitutive signaling of CXCR2 leads to transforming activity similar to Kaposi's sarcoma herpesvirus-G protein-Biochemistry 38(27): 8703-12.

Burstein, E. S., T. A. Spalding, et al. (1996). "Constitutive activation of chimeric m2/m5 muscarinic receptors and delineation of G-protein coupling selectivity domains." coupled receptor." Limmunol 163(4): 2017-22.

Cavalli, A., A. M. Babey, et al. (1999). "Altered adenylyl cyclase responsiveness subsequent to point mutations of Asp 128 in the third transmembrane domain of the delta-opioid Biochem Pharmacol 51(4): 539-44.

Charpentier, S., K. R. Jarvie, et al. (1996). "Silencing of the constitutive activity of the dopamine D1B receptor. Reciprocal mutations between D1 receptor subtypes delineate residues underlying activation properties." I Biol Chem 271(45): 28071-6. receptor." Neuroscience 93(3): 1025-31.

Cho, W., L. P. Taylor, et al. (1996). "Mutagenesis of residues adjacent to transmembrane prolines alters D1 dopamine receptor binding and signal transduction." Mol Pharmacol

Cohen, D. P., C. N. Thaw, et al. (1997). "Human calcitonin receptors exhibit agonist-independent (constitutive) signaling activity." <u>Endocrinology</u> 138(4): 1400-5. Cohen, G. B., T. Yang, et al. (1993). "Constitutive activation of opsin: influence of charge at position 134 and size at position 296." <u>Biochemistry</u> 32(23): 6111-5. Dube, P., A. DeCostanzo, et al. (2000), "Interaction between transmembrane domains five and six of the alpha -factor receptor." <u>Unic Chem</u> 275(34): 26492-9.

Duprez, L., J. Parma, et al. (1994). "Germline mutations in the thyrotropin receptor gene cause non- autoimmune autosomal dominant hyperthyroidism." Nat Genet 7(3): 396-401. Egan, C. T., K. Herrick-Davis, et al. (1998). "Creation of a constitutively activated state of the 5- hydroxytryptamine2A receptor by site-directed mutagenesis: inverse ag nist

activity of antipsychotic drugs." I Pharmacol Exp. Ther 286(1): 85-90.

Esapa, C. T., L. Duprez, et al. (1999). "A novel thyrotropin receptor mutation in an infant with severe thyrotoxicosis." Thyroid 9(10): 1005-10.

Fanelli, F., P. Barbier, et al. (1999). "Activation mechanism of human oxytocin receptor: a combined study of experimental and computer-simulated mutagenesis." Mol Pharmagol

Gaudin, P., J. J. Maoret, et al. (1998). "Constitutive activation of the human vasoactive intestinal peptide 1 receptor, a member of the new class II family of G protein-coupled receptors." J Biol Chem 273(9): 4990-6.

Gaudin, P., C. Rouyer-Fessard, et al. (1998). "Constitutive activation f the human VIP1 receptor." Ann N. Y. Acad Sci 865: 382-5.

Groblewski, T., B. Maigret, et al. (1997). "Mutation of Asn 111 in the third transmembrane domain of the AT1A angiotensin II receptor induces its constitutive activation." I Biol FIG. 1 (14 of 15)
Govardhan, C. P. and D. D. Oprian (1994). "Active site-directed inactivation of constitutively active mutants of rhodopsin." I Biol Chem 269(9): 6524-7.

Han, M., S. O. Smith, et al. (1998). "Constitutive activation of opsin by mutation of methionine 257 on transmembrane helix 6." Biochemistry 37(22): 8253-61

Hasegawa, H., M. Negishi, et al. (1996). "Two isoforms of the prostaglandin E receptor EP3 subtype different in agonist-independent constitutive activity." I Biol Chem 271(4):

Herrick-Davis, K., C. Egan, et al. (1997). "Activating mutations of the serotonin 5-HT2C receptor." INsurocham 69(3): 1138-44.

Hjorth, S. A., C. Orskov, et al. (1998). "Constitutive activity of glucagon receptor mutants." Mol Endoctinal 12(1): 78-86.

Högger, P., M. S. Shockley, et al. (1995). "Activating and mactivating mutations in N. and C-terminal 13 loop junctions of muscarinic acetylcholine Hm1 receptors." J Biol Chem

Hwa, J., R. Gaivin, et al. (1997). "Synergism of constitutive activity in alpha 1-adrenergic receptor activation." <u>Biochemistry</u> 36(3): 633-9.
Ishii, I., T. Izumi, et al. (1997). "Alanine exchanges of polar amino acids in the transmembrane domains of a platelet-activating factor receptor generate both constitutively active and inactive mutants." I Biol Chem 272(12): 7846-54.

Jensen, A. A., T. A. Spalding, et al. (2000). "Functional importance of the Alal 16-Pro136 region in the calcium-sensing receptor. CONSTITUTIVE ACTIVITY AND INVERSE AGONISM IN A FAMILY C G-PROTEIN-COUPLED RECEPTOR [In Process Citation]." I Biol Chem 275(38): 29547-55.

Jin, J., G. F. Mao, et al. (1997). "Constitutive activity of human prostaglandin B receptor BP3 isoforms." British I Pharmacol 121: 317-23.

Kjelsberg, M. A., S. Cotecchia, et al. (1992). "Constitutive activation of the alpha 1B-adrenergic receptor by all amino acid substitutions at a single site. Evidence for a region which constrains receptor activation." I Biol Chem 267(3): 1430-3.

Konopka, J. B., S. M. Margarit, et al. (1996). "Mutation of Pro-258 in transmembrane domain 6 constitutively activates the G protein-coupled alpha-factor receptor." Proc Nati Acad Sci U S A 93(13): 6764-9

Kosugi, S., C. Van Dop, et al. (1995). "Characterization of heterogeneous mutations causing constitutive activation of the luteinizing hormone receptor in familial male precocious puberty." Hum Mol Genet 4(2): 183-8.

Kudo, M., Y. Osuga, et al. (1996). "Transmembrane regions V and VI of the human luteinizing hormone receptor are required for constitutive activation by a mutation in the third intracellular loop." J Biol Chem 271(37): 22470-8.

Lefkowitz, R. J., S. Cotecchia, et al. (1993). "Constitutive activity of receptors coupled to guanine nucleotide regulatory proteins." Trands Pharmacol Sci 14(8): 303-7.

Liu, J., N. Blin, et al. (1996). "Molecular mechanisms involved in muscarinic acetylcholine receptor- mediated G protein activation studied by insertion mutagenesis." J Biol Chem 271(11): 6172-8.

Marie, J., C. Koch, et al. (1999). "Constitutive activation of the human bradykinin B2 receptor induced by mutations in transmembrane helices III and VI." Mol Pharmacol 55(1):

Mason, D. A., J. D. Moore, et al. (1999). "A gain-of-function polymorphism in a G-protein coupling domain of the human beta l-adrenergic receptor." I Biol Chem 274(18):

Matus-Leibovitch, N., D. R. Nussenzveig, et al. (1995). "Truncation of the thyrotropin-releasing hormone receptor carboxyl tail causes constitutive activity and leads to impaired responsiveness in Xenopus oocytes and A(T20 cells." I Biol Chem 270(3): 1041-7.

Morin, D., N. Cotte, et al. (1998). "The D136A mutation of the V2 vasopressin receptor induces a constitutive activity which permits discrimination between antagonists with partial agonist and inverse agonist activities." FEBS Lett 441(3): 470-5.

Nanevicz, T., L. Wang, et al. (1996). "Thrombin receptor activating mulations. Alteration of an extracellular agonist recognition domain causes constitutive signaling." <u>I Biol</u>

Olesnicky, N. S., A. J. Brown, et al. (1999). "A constitutively active G-protein-coupled receptor causes mating self- compatibility in the mushroom Coprinus." Embo 1 18(10):

Parent, J. L., C. Le Gouill, et al. (1996). "Mutations of two adjacent amino acids generate inactive and constitutively active forms of the human platelet-activating factor receptor." Biol Chem 271(14): 7949-55.

## FIG. 1 (15 of 15)

Parnot, C., S. Bardin, et al. (2000). "Systematic identification of mutations that constitutively activate the angiotensin II type 1A receptor by screening a randomly mutated cDNA Parma, J., J. Van Sande, et al. (1995). "Somatic mutations causing constitutive activity of the thyrotropin receptor are the major cause of hyperfunctioning thyroid adenomas: identification of additional mutations activating both the cyclic adenosine 3,5-monophosphate and inositol phosphate-Ca2+ cascades." Mol Endocrinol 9(6): 725-33 library with an riginal pharmacological bioassay." Proc Natl Acad Sci USA 97(13): 7615-20.

Paschke, R., M. Tonacchera, et al. (1994). "Identification and functional characterization of two new somatic mutations causing constitutive activation of the thyrotropin receptor

Pauwels, P. J., A. Gouble, et al. (1999). "Activation of constitutive 5-hydroxytryptamine B receptor by a series of mutations in the BBXXB motif: positioning of the third intracellular loop distal junction and its goalpha protein interactions [In Process Citation]." Biochem J 343 Pt 2: 435-42. in hyperfunctioning autonomous adenomas of the thyroid." I Clin Endocrinol Metab 79(6): 1785-9.

Perez, D. M., J. Hwa, et al. (1996). "Constitutive activation of a single effector pathway: evidence for multiple activation states of a G protein-coupled receptor." Mol Pharmacol 49(1): 112-22.

Ren, Q., H. Kurose, et al. (1993). "Constitutively active mutants of the alpha 2-adrenergic receptor [published erratum appears in J Biol Chem 1994 Jan 14;269(2):1566]." J Biol Chem 268(22): 16483-7.

Robbins, L. S., J. H. Nadeau, et al. (1993). "Pigmentation phenotypes of variant extension locus alleles result from point mutations that alter MSH receptor function." Cell 72(6): Rim, J. and D. D. Oprian (1995). "Constitutive activation of opsin: interaction of mutants with rhodopsin kinase and arrestin." Biochemistry 34(37): 11938-45.

15/27

Russo, D., M. G. Wong, et al. (1999). "A Val 677 activating mutation of the thyrotropin receptor in a Hurthle cell thyroid carcinoma associated with thyrotoxicosis." Thyroid 9(1):

Samama, P., S. Cotecchia, et al. (1993). "A mutation-induced activated state of the beta 2-adrenergic receptor. Extending the ternary complex model." Journal of Biological

Chemistry 168(7): 4625-36.
Scheer, A., T. Costa, et al. (2000). "Mutational analysis of the highly conserved arginine within the Glu/Asp-Arg-Tyr motif of the alpha(1b)-adrenergic receptor: effects on receptor isomerization and activation." Mol Pharmacol 57(2): 219-31.

Scheer, A., P. Fancili, et al. (1997). "The activation process of the alphal B-adrenergic receptor: potential role of protonation and hydrophobicity of a highly conserved aspartate."


Proc Natl Acad Sci USA 94(3): 808-13. Schipani, E., G. S. Jensen, et al. (1997). "Constitutive activation of the cyclic adenosine 3', 5'-monophosphate signaling pathway by parathyroid hormone (PTH)/PTH-related Shenker, A., L. Laue, et al. (1993). "A constitutively activating mutation of the luteinizing hormone receptor in familial male precocious puberty [see comments]." Nature peptide receptors mutated at the two loci for Jansen's metaphyseal chondrodysplasia." Mol Endocrinol 11(7): 851-8.

Sommers, C. M., N. P. Martin, et al. (2000). "A limited spectrum of mutations causes constitutive activation of the yeast alpha-factor receptor." Biochemistry 39(23): 6898-909.


Spalding, T. A., E. S. Burstein, et al. (1998). "Identification of a ligand-dependent switch within a muscarinic receptor." J Biol Chem 273(34): 21563-8.
Spalding, T. A., E. S. Burstein, et al. (1997). "Constitutive activation of the m5 muscarinic receptor by a series of mutations at the extracellular end of transmembrane 6." Biochemistry 36(33): 10109-16.


Tseng, C. C. and L. Lin (1997). "A point mutation in the glucose-dependent insulinotropic peptide receptor confers constitutive activity." Biochem Biophys Res Commun 232(1):

Wonerow, P., T. Sch neberg, et al. (1998). "Deletions in the third intracellular loop of the thyrotropin receptor. A new mechanism for constitutive activation." LBiol Chem 273(14): 7900-5.



and the first term that the second of the se





A Point Mutation Confers Constitutive Activity to the Rat  $\mu$  Opioid Receptor N=11 Wild type <u>11</u> 150K ¬ 100K -50K -0 FIG. 6 Light Units (CPS) (SRE-luc + Gq5i)

Mutant

20/27

 $\square$  basal  $l \mu$  opiod

pcDNA1

1 µM DAMGO

FIG 7

Target Residues Within Class I GPCR's






FIG. 8



о С П









FIG. 11



| FIG.         | 12                   | 26/27  |
|--------------|----------------------|--|
| erk          |                      | mesi prgepgetcaesactppessawppgwarf ingsaeseeac   |
| orkr         | 3                    | MEST OFFICE SEAMER AND THE PROPERTY OF THE PRO |
| orm          | 1 M                  | MDSSAAPTNASNCTDAEAYSSCSPAPSEGSWUNLSHLDENLSEPCGPNRTDLEGRDSL   |
| ormr<br>ord  | 1 -                  | MDSSTGPGNTSDCSDPEAQASCSPAPGSWENLSHVDCNQSDPCGLNRTGLGGNDSL<br>MEPAPSAGAERO.PPLFENASDAYPSACPSAGANASG  |
| ATla         | 1 -                  | MALNSSAED@IKRI@  |
| BK-2         | 1 -                  | mfspwkismflsvredsvptt <b>a</b> sfs <b>ad</b> mlnvtlogptln <mark>e</mark> .tfao   |
| ork 🐺        | 49 I                 | LEPARISPARIPVERITANYSMVEVVGLMGNSLVMHVTHRYTKMKTATNIYLENLALADA   |
| orkr         | 49 I                 | LEPARISPAI PVIITAMYSMYEVUGIMGNSLVMHVIMRYTKVKTAINIYIENLALADA<br>LEPARISPAI PVIITAMYSMYEVUGIMGNSLVMHVIHRYTKVKTAINIYIENLALADA<br>CPPTGS . PSMITAITIMALYSMYCVVGIFGNELVMMVIMRYTKVKTAINIYIENLALADA   |
| orm          | 59 C                 | CPQTGS.PSMITAITIMALYSIIVCVVGIFGMFLVVKVVIQEYTKVKVKVXINIYIFMLALADA   |
| ord          | 37 1                 | PPGARSASSIVALAITAN SAWCAVERSON NAVINGO IN TO VINCE SAVING TO THE PROPERTY OF T |
| ATla         | 16 I                 | DDCPRAGRHSYI FWW PPT PASIE TAYER FONSIAWI VIYFYM CHAVASWYLL MANADL   |
| EK-2         | 45 5                 | SKCPOVEWLGWLNTHOPPFLWVFDVFATTENIFVFSVFCLHKSSCHVAETVIGNLAAADL   |
| ork          | 107                  | LVTIITIN PEOSTVYLMN . SWPEGINLCKIVISIDYXNWETSIETLIN MSVDRYIAVCHPVK   |
| orkr<br>orm  | 107                  | WHITE PROSERVE WE TRUBERTH OF DUST DYVINETS FOR COMSUDEY AVERPUR   |
| ormr         | 116                  | LVTHTMPFQSAVYLMN.SWPFGDYLCKIVISIDYYNWFTSIFTLTHMSVDRYIAVCHPVK<br>LATSTLPFQSYNYLMG.HWPFGTHLCKIVISIDYYNWFTSIFTLCTMSVDRYIAVCHPVK<br>LATSTLPFQSYNYLMG.HWPFGTHLCKIVISIDYYNWFTSIFTLCTMSVDRYIAVCHPVK   |
| ord          | 97 [                 | LATETINPFOSAKYLME. IMPEGET LCKAVISIDYYNMFTS LFTLTMMSVDRYLAVCHPVK<br>CFLLTMPLWAYYTAMEYRWDFGNHLCKIASASVTENIYASMFLLMCISIDRYHAIVHPMK   |
| ATLA<br>BK-2 | 76 (<br>105 <b>!</b> | CFILING LWAYYTAMEYRWIGGHILGCIASESVIEWIYASMILLACUSEDRYDALVKIYS<br>ILACGIAPEWATISNNFDWLEGETICEVVNETISMWAYESICFLWLYSEDRYDALVKIYS  |
|              | -                    | -14 from DRY   |
| ork          | 166                  | ALDERTELKAKI INI CIWELSSYGI SARVLEGTKVR. EDVDVI ECSLOFEDDDYSWD<br>ALDERTELKAKI INI CIWELASSYGI SARVLEGTKVR. EDVDVI ECSLOFEDDEYSWD<br>ALDERTERNAKI INFONWELSSALGI EVWEWATTKYR. O. GS ID CELTSHPTW. WE<br>ALDERTERNAKI WAYONWELSSALGI EVWEWATTKYR. O. GS ID CELTSHPTW. WE  |
| orkr         | 166<br>177           | AMDERIPLKAKOTANI CIMPILASSI GITSAINA GITKUR. BDVD (1200-2140-220DE) SAMO   |
| ormr         | 175                  | AND STANKE TO MENT SEATON PARTY THEY SEATON TO SHE CONTRIBUTE SHOTE  |
| ord :-       | 156 !                | [Jana] 4 (4 h 4 d 1 for 1 and 1 d 1 d 1 d 1 d 1 d 2 d 2 d 2 d 2 d 2  |
| ATla<br>BK-2 | 136                  | SRLRRIMLVAKVICIIIWWWAGLASTPAVIHRNVYFIENINTIVOAFHYESRN.SILP<br>MGRMRGVRWAKYSLVIWGCTLLISSPWAVFRIMKEYSDEGHNVIACVISYPSLIWE   |
| DI. Z        |                      |  |
| ork          | 224                  | LEMKICVFIFAFYIPVLIIMVCYTLMILRLKSVRILSGSREKDRNLRRITRLVLVVVAVF<br>LEMKICVFMFAFYIPVLIIMVCYTLMILRLKSVRILSGSREKDRNLRRITRUVLVVVAVF<br>NLIKICVFIFAFIMPVLIIMVCYGLMILRLKSVRIMLSGSKEKDRNLRRITRUVLVVVAVF<br>NLIKICVFIFAFIMPVLIIMVCYGLMILRLKSVRIMLSGSKEKDRNLRRITRUVLVVVAVF   |
| orkr<br>orm  | 224                  | ILPMKTCVFMFAFYTPVLITTUVCYTLMTLRLKSVRJLSGSREKDRNLRRTTRUVLVVAVE<br>NUKTOVFTFAFTMPVLTTOVCYCLMTTRLKSVRMLSGSKEKDRNLRRTTRUVLVVVAVE   |
| ormr         | 230                  | NLCKICVFIFAFILIPVLIIIVCYGLMILRLKSVRVLSGSKEKDRNLRRITRNVLVVVAVE  |
| ord          | 211                  | TVT::eleviside/bitationnial/TVO/GROBING SERVICEINSGESCOCOCOCOCOCOCOCOCOCOCOCOCOCOCOCOCOCOCO  |
| ATla<br>BK-2 | 193<br>222           | igigetknilgetepentetsynniwkalkkayeioknkprnddifraimalivlee<br>vetnwlinivgemie.isviteciwoiwovlennewokekeiote.rraivúvlvvílde  |
|              |                      |  |
| ork          | 284                  | yvcwtpihifilivealgs.tshstaalssmyscialgytnssinpelyafidene<br>tiicwtpihifilivealgs.tshstavilssmyscialgytnssimpvlyafidene<br>tvcwtpihifydeikalytipshiffotvsmhscialgytnscinpvlyafidene   |
| orkr         | 284                  | TUCOUPIE THE ALCS.TSHETAVLS MYTCLALGY INSSEMPLY AFLDENE  |
| orm          | 290                  | Participancy as a style mark which is a second of the following the following second s |
| ord          | 271                  | FINE (OF EAST AND STANKE OF THE TALL OF THE TARKET AND THE TARKET  |
| ATla         | 250                  | FFS@VPHOLETFUDVLIOLGVIHDCKISDIVDT&MPITICLEYFNMCLMPRFYGELGKKE   |
| BK-2         | 280                  | EINOMER POR STEIN TEHREGILS SCODER INDVITQUAS FERREN SEAMOND VINGER BEING SEQ ID NO:   |
| ork          | 778                  |  |
| orkr         | 338                  | KRCERD FOF PEKMRMER OSTSRAR. NIVOD. PAYLRDIDGMNKPV 76 KRCERD FOF PEKMRMER OSTNRAR. NIVOD. PASMRDVGGMNKPV 77  |
| orm          | 346                  | KRCHRESTINION ROMESTANTYDRING MEDICAL TO THE STANTYDRING MEDICAL TO THE STA |
| ormr         | 344                  | KRCERESCIPTSSNIE CONSURIERONY RDHESTANT DRIVED LENLEAETAPLP 78 KRCERESCIPTSSTIE CONSURVIRONY REHESTANT DRIVED LENLEAETAPLP 79 KRCEROLORKPCGRPDPSSFSRAREATARERVTACTPSDGPGGGAAA  |
| ord<br>ATla  | 310                  | 「WYTELOLLKYTPPKAKSHELSLSTKELSTLEYNPSDNESSSAKKPASCFEVE- 81  |
| BK-2         | 340                  | RKKSWEVYQGVCQKGGCRSEPIQMENSKGTLRTSISWERQIHKLQDWAGSRQ 82  |
|              |                      |  |

| FIG. 13   |  | 27/27  |
|---|--|--|
| mCRmouse mORrat mORbovin mORhuman mORpig mORws ATla BK-2                      | 1  | MDSSAGEGNISDCSDPIA.FASCSPA.ESWUNLSHVDGNOSDPOGENEYGLGGSKSLO MDSSTGEGNTSDCSDPIA.OASCSPA.EGSWUNLSHVDGNOSDPOGENEYGLGGSKSLO MDSGAVETNASNCUDEFTHESGCSPAPSESWUNFSHLEGMLSDPOGENEYGLGGSDFLO MDSSAAETNASNCUDATAY.SSCSPAPSESWUNFSHLEGMLSDPOGENEYDLGGSDSLO MDSSADERNASNCUDEFSPSSMCSEVPSHESWUNFSHLEGMLSDPOGENEYDLGGSDSLO MDSSADERNASNCUDEFSPSSMCSEVPSHESWUNFSHLEGMLSDPOTRNEYELGGSDSLO METSGNISDFLYPISNEVMSNSSVLCRNFSNSTSFLNMNGSSRDSTD ——MALNSSAEDGIKRIODDC ——MFSEWKISMFLSVREDSVPTTASFSADMLNVTLOGETING.TFACSKC   |
| mORmouse mORrat mORbovin mORhuman mORpig mORws ATla BK-2                      | 61<br>60<br>61<br>48                                 | POTGSPSWITALTIMALYS IVCVVGLEGNELVWYVIVRYTKVKTATNIYIENLALADALA<br>POTGSPSWITALTIMALYS IVCVVGLEGNELVWYVIVRYTKVKTATNIYIENLALADALA<br>PSAGSPSWITALTIMALYS IVCVVGLEGNELVWYVIVRYTKVKTATNIYIENLALADALA<br>PPTGSPSWITALTIMALYS IVCVVGLEGNELWWYVIVRYTKVKTATNIYIENLALADALA<br>PPTGSPSWITALTIMALYS IVCVVGLEGNELWWYVIVRYTKVKTATNIYIENLALADALA<br>PPTGSPSWITALTIMALYS IVCVVGLEGNELWWYVIVRYTKVKTATNIYIENLALADALA<br>PKAGRHSYIEVW. IPTIMSITTEVVGTEGNELWWIVITRYTKVKTATNIYIENLALADALA<br>PKAGRHSYIEVW. IPTIMSITTEVVGTEGNELWWIVIVYYLERYTKVKTVASWELLALADALA<br>PKAGRHSYIEVW. IPTIMSITTEVVGTEGNELWWIVIVYYLERYTKVKTVASWELLALADALGE<br>POVEWLGWENTT. OPPFLWWIEVWTTLENIEVTSWFCLHKSSOTVAEIYTGNLAAADLIL |
| mORmouse mORrat mORbovin mORhuman mORpig mORws AT1a BK-2                      | 118<br>118<br>121<br>120<br>121<br>107<br>78<br>107  | TSTLPFQSVNYLMG. TWPFGNILCKIVISIDYYMFTSIFTLCTMSVDRYLAVCHPVKAL TSTLPFQSVNYLMG. TWPFGTILCKIVISIDYYMMFTSIFTLCTMSVDRYLAVCHPVKAL TSTLPFQSVNYLMG. TWPFGTILCKIVISIDYYMMFTSIFTLCTMSVDRYLAVCHPVKAL TSTLPFQSVNYLMG. TWPFGTILCKIVISIDYYMMFTSIFTLCTMSVDRYLAVCHPVKAL TSTLPFQSVNYLMG. TWPFGTILCKIVISIDYYMMFTSIFTLCTMSVDRYLAVCHPVKAL TSTLPFQSVNYLMG. TWPFGDVGCLVMSIDYYMMFTSIFTLTTMSÄDRYLAVCHPVKAL TSTLPFQSVNYLMG. TWPFGDVGCLVMSIDYYMMFTSIFTLTTMSÄDRYLAVCHPVKAL TMTLPLWSVYTAMEYRWPFGNHLCKIASASVTENTVASVETLTGTSTDRYTATVHRMKSR ACGLPFWATTISNNFDWLFGETLCTWVNTIISMNLYSSICFLMLVSTDRYTATVRMG  |
| mORmouse<br>mORrat<br>mORbovin<br>mORhuman<br>mORpig<br>mORws<br>AT1a<br>BK-2 | 177<br>177<br>180<br>179<br>180<br>166<br>138<br>167 | DERTPRNAKI MNYCHWILSSAIGLPVMEMATTKYRC GSIDCTLTESHPTWYWE DERTPRNAKI MNYCHWILSSAIGLPVMEMATTKYRC GSIDCTLTESHPTWYWE DERTPRNAKI MECHWILSSAIGLPVMEMATTKYRC GSIDCTLTESHPTWYWE DERTPRNAKI MNYCHWILSSAIGLPVMEMATTKYRC GSIDCTLTESHPTWYWE DERTPRNAKI MNYCHWILSSAIGLPVMEMATTKYRK GSIDCALTESHPTWYWE DERTPRNAKI MNYCHWILSSAIGLPVMEMATTKYRK GSIDCALTESHPTWYWE DERTPRNAKI MNYCHWILSSAIGLPVMEMASTTIENONSPLOVSNEDCTLLEPHEPWYWE LREIMLVAKOTCIII WEMAGLASLEAVIHRNV YFIENTNITVCAFHYESRNSTLP RMRGVRWAKIYSLVIWGCTLLISSPMINFRIMK EYSDEGHNVTACVISYPS LIWE   |
| mORmouse<br>mORrat<br>mORbovin<br>mORhuman<br>mORpig<br>mORws<br>AT1a<br>BK-2 | 226<br>193   | NLLKI CVFI FAPIMPVLI ITVCYGLMILRLKSVRMLSGSKEKDRNLRRITRMVLVVVAVF NLLKI CVFI FAPIMPVLI ITVCYGLMI LRLKSVRMLSGSKEKDRNLRRITRMVLVVVAVF NLLKI CVFI FAPIMPJLI ITVCYGLMI LRLKSVRMLSGSKEKDRNLRRITRMVLVVVAVF NLLKI CVFI FAPIMPVLI ITVCYGLMI LRLKSVRMLSGSKEKDRNLRRITRMVLVVVAVF NLLKI CVFI FAPIMPVLI ITVCYGLMI LRLKSVRMLSGSKEKDRNLRRITRMVLVVVAVF TLLKI CVFI DAPIMPVLI ITVCYGLMI LRLKSVRMLSGSKEKDRNLRRITRMVLVVVAVF IGLGTTKNILGFTSPFLI ILTSYTLIMKALKKAYETOKNKRRITRMVLVVVAVF VFTNALINVVGFTLF. LSVITFCTMOIMOVLRNNDVOKFKETOTE. RRATVLVLVVL   |
| mORmouse<br>mORrat<br>mORbovin<br>mORhuman<br>mORpig<br>mORws<br>ATla<br>BK-2 | 290<br>293<br>292<br>293<br>286<br>250               | IVCWTPIHIYVIIKALŪTI PETTFQTVSWHFCIALGYTNSCLNPVLYAFLDENFIVCWTPIHIYVIIKALITI PETTFQTVSWHFCIALGYTNSCLNPVLYAFLDENF   |
| mORmouse<br>mORrat<br>mORbovin<br>mORhuman<br>mORpig<br>mORws<br>AT1a<br>EK-2 | 344<br>344<br>347<br>346<br>340<br>310               | KRCFREFC TPTSSTTEQONSTRÜRONTREHPSTANTVORTNHOLENLEAETAPLE 79 KRCFREFC TPTSSTTEQONSTRÜRONTREHPSTANTVORTNHOLENLEAETAPLE 84 KRCFREFC TPTSSNTEQONSTRÜRONTREHPSTANTVORTNHOLENLEAETAPLE 85 KRCFREFC TPTSSTTEQONSARTRONTREHPSTANTVORTNHOLENLEAETAPLE 86 KRCFREFC PSSPSVLELONSTRUSNPOQESQESCHKVDRUNGOV 87 KKYFLOLLKYTEPKÄKSHS SLSTKMSTLSYRPSDAMSSSAKKPASCFEVE 81  |